

**Purdue University**  
**Purdue e-Pubs**

---

Open Access Theses

Theses and Dissertations

---

January 2016

# Effect of Obesity on the Skeletal Muscle Angiogenic Response Following Acute Resistance Exercise

Jessica Aryn Weiss  
*Purdue University*

Follow this and additional works at: [https://docs.lib.purdue.edu/open\\_access\\_theses](https://docs.lib.purdue.edu/open_access_theses)

---

## Recommended Citation

Weiss, Jessica Aryn, "Effect of Obesity on the Skeletal Muscle Angiogenic Response Following Acute Resistance Exercise" (2016).  
*Open Access Theses*. 1137.  
[https://docs.lib.purdue.edu/open\\_access\\_theses/1137](https://docs.lib.purdue.edu/open_access_theses/1137)

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact [epubs@purdue.edu](mailto:epubs@purdue.edu) for additional information.

**PURDUE UNIVERSITY**  
**GRADUATE SCHOOL**  
**Thesis/Dissertation Acceptance**

This is to certify that the thesis/dissertation

prepared By Jessica Aryn Weiss

Entitled

EFFECT OF OBESITY ON THE SKELETAL MUSCLE ANGIOGENIC RESPONSE FOLLOWING ACUTE RESISTANCE EXERCISE

For the degree of Master of Science

Is approved by the final examining committee:

Timothy Gavin

Chair

Bruno Tesini Roseguini

Tara M. Henagan

To the best of my knowledge and as understood by the student in the Thesis/Dissertation Agreement, Publication Delay, and Certification Disclaimer (Graduate School Form 32), this thesis/dissertation adheres to the provisions of Purdue University's "Policy of Integrity in Research" and the use of copyright material.

Approved by Major Professor(s): Timothy Gavin

Approved by: David B. Klenosky

Head of the Departmental Graduate Program

6/30/2016

Date

EFFECT OF OBESITY ON THE SKELETAL MUSCLE ANGIOGENIC RESPONSE  
FOLLOWING ACUTE RESISTANCE EXERCISE

A Thesis

Submitted to the Faculty

of

Purdue University

by

Jessica A Weiss

In Partial Fulfillment of the

Requirements for the Degree

of

Master of Science

August 2016

Purdue University

West Lafayette, Indiana

## TABLE OF CONTENTS

	Page
LIST OF FIGURES .....	iv
LIST OF TABLES .....	v
LIST OF ABBREVIATIONS .....	vi
ABSTRACT .....	viii
CHAPTER 1. INTRODUCTION .....	1
1.1 Obesity and Insulin Resistance .....	1
1.2 The Impact of Exercise .....	2
1.3 Skeletal Muscle Angiogenesis .....	3
1.4 Obesity .....	7
1.5 Exercise and Angiogenesis .....	8
1.6 Obesity, Exercise, & Angiogenesis .....	9
1.7 Hypothesis .....	10
CHAPTER 2. METHODS .....	12
2.1 Participant Selection .....	12
2.2 Experimental Design .....	13
2.3 Statistical Analysis .....	18
CHAPTER 3. RESULTS .....	19
3.1 Subject Characteristics .....	19
3.2 VEGF and VEGF receptor expression .....	19
3.3 TSP-1 expression .....	21
3.4 AMPK and p-AMPK expression .....	22
CHAPTER 4. DISCUSSION .....	24
4.1 Obesity and basal VEGF and VEGF receptor expression .....	24

	Page
4.2 Obesity, exercise and VEGF and VEGF receptor expression .....	25
4.3 Obesity and basal TSP-1 expression .....	26
4.4 Obesity, exercise and TSP-1 expression .....	28
4.5 Obesity and basal p-AMPK and AMPK expression .....	28
4.6 Obesity, exercise and p-AMPK and AMPK expression .....	29
4.7 Limitations.....	31
4.8 Future Directions .....	32
CHAPTER 5. CONCLUSION .....	33
REFERENCES .....	34

## LIST OF FIGURES

Figure	Page
Figure 1: Angiogenic Pathway in Response to Exercise .....	7
Figure 2: Day 2 timeline .....	15
Figure 3. Skeletal muscle vascular endothelial growth factor (VEGF) .....	20
Figure 4. Skeletal muscle vascular endothelial growth factor receptor (VEGFR2). .....	21
Figure 5. Skeletal muscle thrombospondin-1 (TSP-1) .....	22
Figure 6. Skeletal muscle 5' AMP-activated protein kinase (AMPK) .....	23

## LIST OF TABLES

Table	Page
Table 1: Subject Characteristics.....	13
Table 2: Primer sequences for gene expression.....	18

## LIST OF ABBREVIATIONS

NHLBI	National Heart, Lung, and Blood Institute
T2DM	Type II diabetes mellitus
NIDDK	National Institute of Diabetes and Digestive and Kidney Diseases
EC	Endothelial cells
VEGF	Vascular endothelial growth factor
eNOS	Endothelial nitric oxide synthase
NO	Nitric oxide
TSP-1	Thrombospondin-1
AMPK	AMP-activated protein kinase
ADP	Adenosine diphosphate
ATP	Adenosine triphosphate
AMP	Adenosine monophosphate
p-AMPK	phosphorylated AMPK
MAPK	Mitogen-activated protein kinase
BMI	Body mass index
LN	Lean
OB	Obese
PAR-Q	Physical activity readiness questionnaire



VO <sub>2</sub> MAX	Maximal oxygen consumption
1RM	1 repetition maximum
RPE	Rate of perceived exertion
RER	Respiratory exchange ratio
HDL	High density lipoprotein
LDL	Low density lipoprotein
FSG	Fasting Serum Glucose
FSI	Fasting Serum Insulin
ANOVA	Analysis of variance

## ABSTRACT

Weiss, Jessica A. M.S., Purdue University, August 2016. Effect of Obesity on the Skeletal Muscle Angiogenic Response Following Acute Resistance Exercise. Major Professor: Timothy Gavin.

With the rates of obesity remaining high, health risks are abundant. This includes an increased possibility of the development of insulin resistance and type 2 diabetes mellitus (T2DM). Several biochemical and histochemical characteristics associated with obesity have been reported, including a reduction in capillary density in the skeletal muscle and increased muscle fiber size. Resistance exercise is known to induce hypertrophy and increase skeletal muscle capillarization. Following acute resistance exercise, it is hypothesized angiogenic factors are reduced and angiostatic factors are increased in the skeletal muscle of obese individuals. Muscle biopsies from the vastus lateralis were obtained to investigate whether VEGF, VEGF receptor, TSP-1 expression, and AMPK phosphorylation (an indicator AMPK activation) are lower in obese compared to lean individuals at rest and in response to an acute bout of exercise. Muscle biopsies were taken before, 15 MIN, and 3 HR following 3 sets of single-leg knee extension at 80% 1 RM. Eight sedentary lean (4 male, 4 female) and eight sedentary obese (4 male, 4 female) individuals participated in this study. No differences in VEGF or VEGF receptor mRNA and protein levels were found at rest or following exercise in lean or obese individuals. Exercise increased VEGF mRNA independent of group. Resistance exercise did not affect TSP-1 mRNA and protein expression or AMPK

phosphorylation and no differences between lean and obese were found. The present study suggests that the angiogenic response to resistance exercise (a known hypertrophic stimulus) is similar in lean and obese individuals.

## CHAPTER 1. INTRODUCTION

### 1.1 Obesity and Insulin Resistance

The prevalence of obesity in the United States remains high, with 78.6 million adults currently falling under this classification (Centers for Disease Control and Prevention, 2014). Many chronic conditions are associated with obesity, which has earned it the title of the second leading cause of preventable death in the United States by the National Heart, Lung, and Blood Institute (NHLBI). Health problems associated with obesity include hypertension, heart disease, metabolic syndrome, and type 2 diabetes mellitus (T2DM) (Mayo Clinic, 2015). In 2008, the estimated annual medical cost of obesity in the United States was \$147 billion and has been projected to rise due to the increasing prevalence of T2DM (Centers for Disease Control and Prevention, 2014). According to the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), T2DM frequently is the result of prolonged insulin resistance or a decreased sensitivity and/or responsiveness to insulin. Insulin resistance is common among the obese population. It is estimated that 24% of obese adults in the United States are insulin resistant while in the obese adolescent population prevalence is estimated at 52% (Lee et al, 2006; McKeown et al, 2004).

The human body consists of roughly 40% skeletal muscle, which is responsible for 75% of insulin-induced glucose disposal (Reaven & Laws, 1999).

Several biochemical and histochemical characteristics within the skeletal muscle have been associated with obesity. Studies have shown fiber type dissimilarities between the skeletal muscles of lean compared to obese individuals. A decrease in Type I, slow twitch oxidative, fibers concurrent with an increase in Type IIB, fast twitch glycolytic, fibers has been noted in obese individuals (Lillioja et al, 1987). The area of skeletal muscle fibers of obese individuals is significantly larger than the skeletal muscle fibers of lean individuals (Gavin et al, 2005). Reduced mitochondrial function, reducing the oxidative capacity of the skeletal muscle, has also been reported in obese individuals (Torgan et al, 1989). Primarily, obesity has been strongly correlated to a reduction in capillary density, or the number of capillaries per unit cross-sectional area of muscle (Gavin et al, 2005; Lash et al, 1989; Lillioja et al, 1987; Shono et al, 1999; Torgan et al, 1989). Capillaries are essential for the delivery of oxygen and other nutrients important for energy production. Increased capillarization ensures adequate diffusion capacity across the skeletal muscle (Hudlicka, Brown & Egginton, 1992). Therefore, alterations in skeletal muscle capillarization may contribute to much of the insulin resistance associated with obesity (Lillioja et al, 1987; Shono et al, 1999).

## 1.2 The Impact of Exercise

Exercise has been proven to be beneficial to overall health in a variety of ways. Those who participate in physical activity for 7 hours or more a week reduce their risk of mortality by 40% (Centers for Disease Control and Prevention, 2015). It has been well-established that endurance training provides many improvements to overall health. A host of effects of aerobic exercise support improvements which include cardiac and skeletal

muscle changes. Adaptations in skeletal muscle include increases in mitochondrial content, respiratory capacity, and capillary density of muscle fibers (Andersen & Henriksson, 1977; Hermansen & Wachtlova, 1971; Holloszy & Coyle, 1984; Yamashita et al, 1993). Regular resistance exercise can also improve cardiovascular, bone, and muscle health and reduce the risks of many chronic diseases (American College of Sports Medicine Position Stand, 1990). Muscle hypertrophy observed with resistance exercise contributes to an increase in muscle strength, basal metabolic rate, and lean body mass, while decreasing body fat composition. Resistance exercise has also been shown to benefit glucose metabolism, lowering basal insulin levels and increasing insulin sensitivity (Pollock & Vincent, 1996). Less recognized as an outcome of resistance exercise is an increase in skeletal muscle capillarization (McCall et al, 1996; Schantz, 1982). Increased capillary density improves the exchange of oxygen, substrates, and metabolites between blood and skeletal muscle (Andersen & Henriksson, 1977; Hermansen & Wachtlova, 1971; Poole, Copp, Ferguson, & Musch, 2013; Tesch, Thorsson & Kaiser, 1984), which is vitally important for individuals with chronic conditions where this exchange has been compromised (Adair & Montani, 2010).

### 1.3 Skeletal Muscle Angiogenesis

Angiogenesis, the process of generating new capillary blood vessels, occurs in response to signals from changes in mechanical stimuli and metabolic demand (Hudlicka, Brown & Egginton, 1992). It involves a complex cascade of events leading to the migration and proliferation of capillary endothelial cells (EC) (Engerman et al, 1967).

The interplay of both angiogenic (positive) and angiostatic (negative) factors contribute to this process.

### *Angiogenic and Angiostatic Factors*

One of the most vital pro-angiogenic factors for basal and exercise-induced capillarization is vascular endothelial growth factor (VEGF). VEGF is a 45 kDa heparin-binding vascular EC mitogen of the micro- and macrovascular EC of arteries, veins, and the lymphatics (Amaral et al, 2001; Dvorak et al, 1979; Ferrara & Davis-Smyth, 1997; Ferrara & Henzel, 1989; Gavin TP, 2009; Olfert et al, 2009; Tang et al, 2004; Wagner et al, 2006). Although several different isoforms exists, VEGF-A plays the most crucial role in exercise-induced skeletal muscle angiogenesis (Shweiki et al, 1992; Waltenberger et al, 1994). It is released into the muscle interstitial space in response to mechanical stimuli and other stress inducing factors (Gavin et al, 2007; Hoffner, Nielsen, Langberg, & Hellsten, 2003; Hoier et al, 2011; Hoier et al, 2012; Hoier et al, 2010; Ouchi, Shibata, & Walsh, 2005), including hypoxia, oxidative stress, growth factors, and cytokines (Ikeda et al, 1995; Kuroki et al, 1996; Takahashi et al, 2002; Wang, Huang, Kazlauskas, & Cavenee, 1999).

Two receptors identified as having a high affinity for VEGF are VEGFR1 (de Vries et al, 1992) and VEGFR2. Expression of both receptors is restricted to the vascular endothelium, where the positive charge of the surface of VEGF facilitates its binding (Ferrara & Keyt, 1997; Keyt et al, 1996; Yamaguchi et al, 1993). VEGFR2 is suggested to be the receptor principally involved in angiogenesis (Millauer et al, 1993; Gavin TP, 2009; Waltenberger et al, 1994). The phosphorylation of VEGFR2 initiates several signaling cascades, resulting in increased vascular permeability and the migration,

proliferation, and survival of EC (Connolly et al, 1989; Koch et al, 2011). One important cascade involves VEGFR2 activation of endothelial nitric oxide synthase (eNOS) to produce nitric oxide (NO) which in turn stimulates EC growth and motility (Ziche et al, 1994). The NO cascade activated by VEGFR2 phosphorylation is inhibited by thrombospondin-1 (TSP-1) (Isenberg et al, 2006).

TSP-1 is a 450 kDa multifunctional homotrimeric matrix glycoprotein that is thought to play an important role in the regulation of angiogenesis (Iruela-Arispe et al, 1995; Lawler, Slayter, & Coligan, 1978). Though TSP-1 has been shown to stimulate the growth and migration of smooth muscle cells (BenEzra, Griffin, Maftzir, & Aharonov, 1993; Lawler, 2000; Nicosia & Tuszynski, 1994). It prevents angiogenesis via inhibition of VEGF release, VEGF signal transduction, and by direct interaction with VEGF (Iruela-Arispe, Bornstein, & Sage, 1991; Lawler & Lawler, 2012). VEGF bioavailability is decreased due to the influence of TSP-1 on the structure of the extracellular matrix (Iruela-Arispe et al, 1995; Lawler, 1986; Chen, Herndon, & Lawler, 2000). TSP-1 binding to receptors CD36 and CD47 inhibits VEGFR2 phosphorylation (Primo et al, 2005), which hinders EC association at the site of development (Bagavandoss & ilks, 1990; Dawson et al, 1997), migration of EC, and stimulates apoptosis (Iruela-Arispe, Bornstein, & Sage, 1991). Further, the uptake and clearance of VEGF from the extracellular space is induced by TSP-1 binding directly to VEGF (Gupta et al, 1999; Greenaway et al, 2007; Lawler & Lawler, 2012). Consistent with TSP-1 being an angiostatic factor, skeletal muscle capillarization is lower in TSP-1 knock-out (KO) compared to wildtype (WT) mice (Malek & Olfert, 2009).

### *Intracellular Regulation*



A key player in the regulation of the responses to angiogenesis-inducing stressors is the heterotrimeric enzyme 5'-AMP-activated protein kinase (AMPK) (Hardie, 2003). When two molecules of adenosine diphosphate (ADP) are used to produce energy in the form of adenosine triphosphate (ATP), the result is the production of one molecule of adenosine monophosphate (AMP) (Kalckar, 1943). In order for a cell to maintain a necessary level of energy for proper function, the ratio of ATP to AMP must remain high. When reductions in the ratio occur, AMPK is activated. Reduced ATP:AMP is caused by disruptions in the production of ATP, hypoxia or ischemia, and/or increases in the consumption of ATP through muscle contraction (Kudo et al, 1995; Marsin et al, 2000; Winder & Hardie, 1996). The phosphorylation of upstream kinases induces AMPK to regulate cellular energy usage (Hawley et al, 1995). AMP regulates the activation of AMPK and inhibits AMPK dephosphorylation by protein phosphatases (Hawley et al, 1995).

The activation of AMPK stimulates energy-conserving/producing pathways to increase while simultaneously decreasing energy-consuming pathways. Previous research suggests AMPK regulates VEGF in the skeletal muscle through the activation of p38 mitogen-activated protein kinase (MAPK) (Pages et al, 2000; Ouchi, Shibata, & Walsh, 2005; Winzen et al, 1999). AMPK also plays a role in the mediation of NO activity (Zhu & Smart, 2005). Increased AMPK activity has been shown to increase VEGF mRNA and VEGF protein, thereby increasing capillarization and the inactivation of AMPK reduces capillarization at the basal level (Ouchi et al, 2005; Zwetsloot, Westerkamp, Holmes & Gavin, 2008). However, AMPK activation is not necessary for aerobic

exercise-induced angiogenesis, but may play a role in response to resistance exercise (Zwetsloot, Westerkamp, Holmes & Gavin, 2008).




Figure 1: Angiogenic Pathway in Response to Exercise.<sup>1</sup>Ouchi, Shibata & Walsh, 2005-*will not be investigated in this report* <sup>2</sup>Ziche et al, 1994; Isenberg et al, 2006-*will not be investigated in this report*

#### 1.4 Obesity

The skeletal muscle fiber size of obese individuals is markedly increased compared to their lean counterparts (Gavin et al, 2005). VEGF is a necessary component for muscle hypertrophy (Huey, Smith, Sulaeman & Breen, 2016), and basal regulation of skeletal muscle VEGF and capillarization is altered by resting levels of AMPK (Zwetsloot, Westerkamp, Holmes, & Gavin, 2008). However, resting levels of metabolic and regulatory factors of angiogenesis are similar between lean and obese individuals of similar physical inactivity. No differences at rest in AMPK, VEGF, or VEGFR2 expression have been found despite reduced capillary density and muscle fiber hypertrophy observed in obese subjects (Gavin et al, 2005; Steinberg et al, 2004). Though comparisons of basal TSP-1 in the skeletal muscle of lean and obese have not been examined, elevated basal TSP-1 protein and gene expression in adipose tissue resulting in

increased circulating TSP-1 in obese subjects has been noted (Abu-Farha, 2013; Hida et al, 2000; Inoue et al, 2013; Varma et al, 2008; Voros et al, 2005).

## 1.5 Exercise and Angiogenesis

### *Aerobic Exercise*

Angiogenesis in response to aerobic exercise in lean subjects has been well researched. The interplay of angiogenic and angiostatic factors lead to an increase in capillary density of the skeletal muscle following endurance training (Andersen & Henriksson, 1977). Studies consistently report elevations in VEGF and VEGFR2 expression in the skeletal muscle in response to acute endurance exercise (Benoit et al, 1999; Gavin et al, 2005; Gustafsson et al, 2005; Ryan et al, 2006). Furthermore, TSP-1 is temporarily increased following acute aerobic exercise, appearing to modulate capillary formation (Hoier et al, 2012; Olfert et al, 2006; Olfert, Wagner, & Power, 2000). Cellular energy usage is regulated through an increase of AMPK activity, heavily dependent upon the duration and intensity of the exercise session (Fujii et al, 2000; Rasmussen & Winder, 1997; Stephens et al, 2002; Winder & Hardie, 1996; Wojtaszewski et al, 2000). AMPK is not a necessary part of the angiogenic response to aerobic exercise (Zwetsloot, Westerkamp, Holmes, & Gavin, 2008) but is one potential regulator of VEGF expression, a key component for the muscle hypertrophic response to resistance exercise (Huey, Smith, Sulaeman & Breen, 2016).

### *Resistance Exercise*

There is very little research regarding the skeletal muscle angiogenic response following resistance exercise. However, it is known that resistance training increases

capillarization in the skeletal muscle in lean subjects (McCall et al, 1996; Schantz, 1982). The increased capillarization is in proportion to the hypertrophy of the muscle fibers in order to maintain the capillary density of the skeletal muscle (McCall et al, 1996; Schantz, 1982). The angiogenic response to resistance exercise in terms of VEGF and VEGFR2 expression is similar to that seen following aerobic exercise (Doyle & Haas, 2010; Huey, Smith, Sulaeman & Breen, 2016; Degens, Moore, & Alway, 2003; Gavin et al, 2007; Parvaresh et al, 2010). Increases in VEGF mRNA are observed 2-3 h post resistance exercise (Gavin et al, 2007; Silvennoinen et al, 2015) and VEGFR2 mRNA 4 h post resistance exercise (Gavin et al, 2007). These increases in VEGF and VEGFR2 expression has been shown to play important roles in the maintenance of capillarization, which is necessary for proper hypertrophy of the muscle (Huey, Smith, Sulaeman & Breen, 2016). Elevated AMPK activity has also been noted in lean individuals during and 1 h following resistance exercise (Dreyer et al, 2006), indicating a potential role in regulating VEGF production following resistance exercise. No investigations into TSP-1 expression in the skeletal muscle following resistance exercise have been published.

## 1.6 Obesity, Exercise, & Angiogenesis

Although no differences reside in basal levels of angiogenic factors between lean and obese, obese demonstrate lower capillarization in the skeletal muscle (Gavin et al, 2005). An important stimulator of the angiogenic process is exercise (McCall et al, 1996; Schantz, 1982). Few have reported any differences in the angiogenic response to aerobic exercise between lean and obese individuals (Gavin et al, 2005; Walton et al, 2015). AMPK activity in obese does not change while lean counterparts show an increase in

AMPK activity following endurance exercise, suggesting an impaired response of AMPK to aerobic exercise (Sriwijitkamol et al, 2007). However, VEGF and VEGFR2 expression in response to aerobic exercise is similar between lean and obese individuals (Gavin et al, 2005). Levels of TSP-1, a potent angiostatic factor, in the adipose tissue of obese individuals return to normal following acute exercise (Abu-Farha et al, 2013).

Although the hypertrophic response to resistance training is evident in obese individuals (Donnelly et al, 1993; Geisler et al, 2011), researchers have not compared the skeletal muscle angiogenic response following resistance exercise in lean and obese individuals to date. The lack of differences in VEGF and VEGFR2 expression following aerobic exercise, combined with the increased fiber diameter with subsequent decreased capillarization (Gavin et al, 2005; McCall et al, 1996), suggests a decreased angiogenic response to muscle hypertrophy in the skeletal muscle of obese individuals.

## 1.7 Hypothesis

At rest, obese subjects will have increased TSP-1 expression compared to lean counterparts. Following acute resistance exercise, obese subjects will have: (i) lower VEGF, VEGF receptor, and AMPK phosphorylation; and (ii) greater TSP-1 expression compared to lean counterparts.

### *Delimitations*

1. Participants were non-smoking and sedentary between the ages 18 and 35. Sedentary specifies participating in physical activity 3 or fewer times per week for 20 minutes or less. One group of individuals were lean with a body mass index (BMI)  $\leq 25 \text{ kg/m}^2$  while the other group was classified as obese with a body mass index (BMI)  $\geq 30 \text{ kg/m}^2$ .

2. Workloads were tailored to each participant by first establishing their 1 repetition maximum (1RM).

## CHAPTER 2. METHODS

The Purdue University Institutional Review Board approved the methods and protocol of this study preceding its execution.

### 2.1 Participant Selection

Eight healthy, sedentary, lean (LN) individuals (4 women; 4 men) and eight healthy, sedentary, obese (OB) individuals (4 women; 4 men) between the ages of 18-35 were recruited to participate in this study. Subjects were nonsmoking and had no known chronic disease in order to be considered healthy enough to participate. Sedentary qualifies as participating in physical activity 3 times or less for 20 minutes or fewer per week. Lean individuals had a body mass index (BMI)  $< 25 \text{ kg/m}^2$ , while obese individuals had a BMI  $\geq 30 \text{ kg/m}^2$ . Qualified individuals were given both verbal and written descriptions of the study. Subjects provided signed consent prior to the beginning of the study, as well as completed an Activity and Tobacco Questionnaire and a Physical Activity Readiness Questionnaire (PAR-Q). Subjects were given instructions on completing a three day food intake diary for the days prior to the Day 2 procedures. Subject characteristics are listed in Table 1.

Table 1: Subject Characteristics. Values are listed as Mean  $\pm$  SE.  $n = 8$  subjects/group.

	Lean	Obese	<i>P</i> Value
Age, yr	21.5 $\pm$ 0.5	24.8 $\pm$ 1.6	0.09
Height, cm	169.6 $\pm$ 2.9	172.8 $\pm$ 3.2	0.48
Mass, kg	62.4 $\pm$ 2.0	109.9 $\pm$ 7.8	<0.01
BMI	21.7 $\pm$ 0.8	36.6 $\pm$ 1.8	<0.01
VO <sub>2</sub> Max, ml O <sub>2</sub> /kg/min	37.3 $\pm$ 2.7	26.1 $\pm$ 2.7	0.01
1 RM, kg	27.8 $\pm$ 0.7	40.3 $\pm$ 4.7	0.03
1 RM, kg/kg	0.45 $\pm$ 0.01	0.37 $\pm$ 0.04	0.17
Fasting glucose, mM	4.8 $\pm$ 0.1	4.9 $\pm$ 0.2	0.54
Fasting insulin, $\mu$ U/ml	11.8 $\pm$ 1.3	28.3 $\pm$ 4.8	0.01
Fasting HOMA-IR	2.46 $\pm$ 0.29	6.38 $\pm$ 1.23	0.01
Fasting HDL, mg/dl	58.4 $\pm$ 7.3	42.9 $\pm$ 5.4	0.08
Fasting LDL, mg/dl	88.6 $\pm$ 12.8	108.6 $\pm$ 5.2	0.11
Fasting Total Cholesterol, mg/dl	162.8 $\pm$ 12.3	180.6 $\pm$ 7.0	0.17
Fasting Triglycerides, mg/dl	79.4 $\pm$ 7.7	131.6 $\pm$ 22.5	0.05

## 2.2 Experimental Design

Day 1: Subjects reported to the Max E. Wastl Human Performance Laboratory where their height, weight, and dominant leg was recorded prior to the completion of a VO<sub>2</sub>MAX test. Subjects were then given a 15 minute rest period before the completion of a one repetition maximum (1 RM) test for single leg, knee extension.

*Maximal Oxygen Consumption (VO<sub>2</sub>MAX)*



Subject  $\text{VO}_{2\text{MAX}}$  was determined two hours post-prandial using a previously described incremental protocol (Kraus et al, 2004) on a cycle ergometer (Lode Excalibur Sport, Groningen, The Netherlands). A warm up was performed at 50 W for 5 minutes on the cycle ergometer. Immediately following, additional workloads were added every 2 minutes by 25 W until volitional fatigue. Expired air was monitored and recorded via open-circuit spirometry (Truemax 2400, Parvo Metrics, Salt Lake City, UT). Additionally, heart rate (model H1, Polar Electro, Lake Success, NY) and rate of perceived exertion (RPE) was continuously monitored and recorded. Subjects were verbally encouraged to continue as long as possible. Criteria for the establishment of  $\text{VO}_{2\text{MAX}}$  included 1) a heart rate of at least 90% of age predicted maximum ( $220 - \text{age}$ ); 2) a respiratory exchange ratio (RER) at or above 1.10; and 3) an identifiable plateau ( $\leq 150$  mL increase) in  $\text{VO}_2$  despite continued increase in workload. At least two of three criteria were met for all subjects.

#### *One Repetition Maximum (1RM)*

Maximal knee extension was randomized between subjects using either the dominant or non-dominant leg. Subjects began in a resting position of  $90^\circ$  and were instructed to fully extend their leg undergoing the exercise using a weight of 13.6 kg. Once parallel to the floor, they returned to the initial position and were given a rest for 1 minute. The exercise was repeated with weight increasing by 4.5 kg until the exercised leg was no longer able to reach full extension. Once this point was attained, the previous weight was determined to be the 1RM.



Figure 2: Day 2 timeline

Day 2: Day 2 occurred at least two weeks after Day 1. This was to avoid any influence from the maximal testing incurred on Day 1 on the subjects' performances on Day 2. Three days preceding Day 2, subjects recorded their food intake and partook in a 12 h fast the night prior.

#### *Blood draw*

On the morning of Day 2, subjects reported to the Max E. Wastl Human Performance Laboratory where a fasting blood draw was obtained. The 5-8 mL blood sample was used for the measurement of glucose, insulin, high density lipoprotein (HDL), low density lipoprotein (LDL), triglycerides, and total cholesterol. It was collected in a serum tube, allowed for clot formation for 30 minutes at room temperature, and then refrigerated until analysis by Mid-America Laboratories. Insulin resistance was measured as HOMA-IR via the following equation:

$$\text{HOMA-IR} = \frac{\text{FSG} * \text{FSI}}{22.5 * 18}$$

Fasting serum glucose (FSG) was used in milligrams per deciliter and fasting serum insulin (FSI) was used in microunits per millimeter (Matthews et al, 1985).

#### *Pre- and post-exercise muscle biopsies*

Immediately following the blood draw, the subjects underwent the procedure to obtain the first (PRE) of three muscle biopsies. All biopsies were obtained from the vastus lateralis: one biopsy before exercise from the non-working leg and two biopsies from the working leg at 15 min post (15 MIN) and 3 h post (3 HR) exercise. The working leg was randomized between non-dominant and dominant leg between subjects. On the working leg, the two biopsy sites were separated by at least 3 cm (Costill, Pearson, & Fink, 1988). Lidocaine (1%) was used as a local anesthetic to obtain the biopsy samples. Tissue samples were immediately placed in liquid nitrogen and stored at -80°C for future use.

#### *Exercise protocol*

Subjects performed three sets of knee extension exercise using their working leg immediately following the first biopsy. Workload was set at 80% of their 1RM determined on Day 1. 8-12 repetitions were performed for the first two sets and subjects were instructed to perform to volitional fatigue for the last set (8-17 repetitions). A two minute rest period was allotted between sets. Preliminary analysis found this protocol to sufficiently fatigue the subjects and completion did not require any outside assistance.

#### *Protein Isolation and Western Blot Analysis*

Muscle biopsy samples were frozen in liquid nitrogen and homogenized in RIPA buffer (50 mM Tris-HCl (pH 7.4), 150 mM NaCl, 2 mM EDTA, 0.1% SDS, 0.1% Triton X-100, and 0.5% Deoxycholate) with protease inhibitor cocktail and phosphatase inhibitor (50 mM NaF and 0.2 mM Na<sub>3</sub>VO<sub>4</sub>). Samples were centrifuged at 4°C for 30 minutes to separate supernatant and insoluble material. The supernatant was used to determine protein concentration by BCA kit (Pierce). Western blot analysis was

performed in order to investigate angiogenic growth factor expression. Angiogenic growth factors include VEGF and VEGF receptor (Gavin et al, 2005); and the anti-angiogenic (angiostatic) factor include TSP-1 (Olfert et al, 2006). AMPK phosphorylation was measured with exercise and obesity as AMPK is an important regulator of VEGF (Zwetsloot, Westerkamp, Holmes, & Gavin, 2008).

Equal amounts of homogenized samples were added to 10% gels. Each membrane was loaded with the 3 HR post exercise sample from subject 4 so that membranes could be compared to one another. Membranes were then run through electrophoresis in running buffer (0.1% SDS, 25 mM Tris, and 190 mM Glycine) at 80 V for 10 minutes, or until proteins reached the end of the stacking gel, and then at 100-140 V for 1-2 hours. Gels were transferred to PVDF membranes in transfer buffer (190 mM glycine 25 mM Tris, and 20% Methanol) for 2 hours at 400 mA. Membranes were stained with Ponceau in order to measure total protein loaded in each sample. After imaging, Ponceau was rinsed from the membranes and then membranes were blocked with 5% non-fat milk dissolved in TBST (25 mM Tris (pH 7.2), 150 mM NaCl, and 0.1% Tween 20) for 1 hour at room temperature. Membranes were incubated with primary antibody (VEGF and VEGFR2, Santa Cruz; AMPK, p-AMPK, and TSP-1 Cell Signaling Technology) in 5% non-fat milk dissolved in TBST at 4°C overnight. Membranes were washed with TBST for 10 minutes 3 times and then incubated in anti-rabbit or anti-mouse immunoglobulin G-horseradish peroxidase (Cell Signaling Technology) or fluorescence (Li-COR) conjugated secondary antibody for 1 hour. Membranes were again washed with TBST for 10 minutes 3 times and signals were visualized by fluorescence or chemiluminescence under FluorChem E system (protein simple).

### *Quantitative real-time PCR*

RNA was isolated from muscle tissue samples using Trizol reagent (Invitrogen). First-strand cDNA was generated by random hexamer primers with MMLV Reverse Transcriptase (Invitrogen). Using a SYBR green PCR kit, the expression level of VEGF, VEGFR2, and TSP-1 was determined by real-time PCR reactions in the Roche LightCycler® 480 System (Roche). The internal control for gene expression was 18s expression. The primer sequences are listed below:

Table 2: Primer sequences for gene expression

Gene Name	Forward (5'-3')	Reverse (5'-3')
VEGF	AGGGCAGAATCATCACGAAGT	AGGGTCTCGATTGGATGGCA
VEGFR2	GTGATCGGAAATGACACTGGAG	CATGTTGGTCACTAACAGAAGCA
TSP-1	AGACTCCGCATCGCAAAGG	TCACCACGTTGTTGTCAAGGG
18s	GGCCCTGTAATTGGAATGAGTC	CCAAGATCCAACACTACGAGCTT

### 2.3 Statistical Analysis

Data were analyzed by use of a two-way (2 groups  $\times$  3 time points) mixed-factorial analysis of variance (ANOVA) test. Following a significant  $F$  ratio, Fisher's LSD *post-hoc* analysis was performed. Significance was established at  $\alpha = 0.05$  level and data reported as Mean  $\pm$  SE.

## CHAPTER 3. RESULTS

### 3.1 Subject Characteristics

Obese (OB) subjects had significantly greater body mass than lean (LN) subjects (OB:  $36.6 \pm 1.8$  kg/m<sup>2</sup>; LN:  $21.7 \pm 0.8$  kg/m<sup>2</sup>). LN subjects demonstrated a significantly higher VO<sub>2Max</sub> than OB (LN:  $37.3 \pm 2.7$  ml O<sub>2</sub>/kg/min; OB:  $26.1 \pm 2.7$  ml O<sub>2</sub>/kg/min). OB demonstrated a higher 1 RM for single leg knee extension (LN:  $27.8 \pm 0.7$  kg; OB:  $40.3 \pm 4.7$  kg); however, when corrected for body mass, the difference was no longer statistically significant ( $P = 0.17$ ). No differences were observed in fasting glucose between groups (LN:  $4.8 \pm 0.1$  mM; OB:  $4.9 \pm 0.2$  mM); however OB demonstrated a higher fasting insulin (LN:  $11.8 \pm 1.3$  μU/ml; OB:  $28.3 \pm 4.8$  μU/ml) and fasting HOMA-IR (LN:  $2.46 \pm 0.29$ ; OB:  $6.38 \pm 1.23$ ).

### 3.2 VEGF and VEGF receptor expression

The VEGF mRNA and protein responses to resistance exercise in LN and OB subjects are displayed in Figure 3. Exercise significantly increased VEGF mRNA similarly at 3 HR in both LN and OB subjects. There was no differences in VEGF protein between LN and OB PRE, 15 MIN or 3 HR post exercise.




Figure 3. Skeletal muscle vascular endothelial growth factor (VEGF) mRNA (a) protein (b) and (c) western blot analysis in response to acute resistance exercise in LN and OB individuals. VEGF mRNA was significantly increased at 3 HR. \*Significantly different from PRE ( $P \leq 0.05$ ). Mean  $\pm$  SE.  $n = 8$  subjects/group.

Skeletal muscle VEGFR2 mRNA and protein responses to exercise in LN and OB individuals are displayed in Figure 4. No differences in the VEGFR2 mRNA or protein were found between LN and OB individuals PRE, 15 MIN or 3 HR post exercise. Exercise did not affect VEGFR2 mRNA or protein in either group.




Figure 4. Skeletal muscle vascular endothelial growth factor receptor (VEGFR2) mRNA (a) protein (b) and western blot analysis (c) in response to acute resistance exercise in LN and OB individuals. Mean  $\pm$  SE.  $n = 8$  subjects/group.

### 3.3 TSP-1 expression

Skeletal muscle TSP-1 mRNA and protein responses to exercise in LN and OB individuals are displayed in Figure 5. There were no differences in TSP-1 mRNA between LN and OB at any time point and no effect of exercise. Similarly, there was no effect of obesity or exercise on TSP-1 protein expression.






Figure 5. Skeletal muscle thrombospondin-1 (TSP-1) mRNA (a) protein (b) and western blot analysis (c) in response to acute resistance exercise in LN and OB individuals.

### 3.4 AMPK and p-AMPK expression

Skeletal muscle AMPK and p-AMPK protein responses to exercise in LN and OB individuals are displayed in Figure 6. Exercise and obesity had no effect on p-AMPK, pan-AMPK, or the p-AMPK/pan-AMPK ratio PRE, 15 MIN or 3 HR post exercise.




Figure 6. Skeletal muscle 5' AMP-activated protein kinase (AMPK) protein (a), phosphorylated AMPK (p-AMPK) protein (b) p-AMPK/AMPK protein (c) and western blot analysis (d) in response to acute resistance exercise in LN and OB individuals. Mean  $\pm$  SE.  $n = 8$  subjects/group.

## CHAPTER 4. DISCUSSION

The principal findings of the current study are there were no differences in the expression of VEGF, VEGF receptor and TSP-1 at rest and following acute resistance exercise between lean and obese groups. Consistent with previous work, resistance exercise did increase VEGF mRNA in both lean and obese individuals. In addition, there was no difference at rest or in response to exercise in AMPK between lean and obese groups and acute resistance exercise did not increase the phosphorylation of AMPK in either group 15 MIN or 3 HR post exercise.

### 4.1 Obesity and basal VEGF and VEGF receptor expression

Similar basal levels of VEGF and VEGF receptor mRNA between the skeletal muscle of lean and obese individuals have been previously reported (Gavin et al, 2005). No there was no difference in the VEGF and VEGF receptor protein content of the skeletal muscle of lean and obese individuals (Gavin et al, 2005). Our data further support the notion that VEGF and VEGF receptor mRNA and protein expression are similar between lean and obese individuals at rest.

#### 4.2 Obesity, exercise and VEGF and VEGF receptor expression

The current report utilizes a single bout of resistance exercise, a stimulus of muscle hypertrophy when repeated, in order to investigate the reduced capillary density observed with basal hypertrophy seen in obese individuals. Investigations using animal models suggest the skeletal muscle of obese have the ability to hypertrophy but have a decreased response to functional overload, a model for resistance exercise, compared to lean (Katta et al, 1985; Paturi et al, 2010). Similarly, Donnelly and colleagues showed an increased hypertrophy with resistance exercise in obese subjects but did not compare these results to lean counterparts (Donnelly et al, 1993). Therefore, this could result in lesser angiogenesis in obese individuals compared to lean due to less activation of hypertrophic pathways.

Very little research regarding the angiogenic response to resistance exercise in lean individuals has been conducted. Previous investigations show elevations in VEGF mRNA as early as 1 h following the completion of acute resistance exercise in lean individuals (Gavin et al 2007; Silvennoinen et al, 2015; Trenerry et al, 2007). Increases in VEGF mRNA >2.5 fold have been reported in human subjects (Gavin et al, 2007). This is relatively similar to what is shown in the present study, demonstrating an increase of ~2.2 fold 3 h post exercise. VEGF protein expression did not change following exercise, and no differences were found between lean and obese subjects. The present findings are in contrast to a previous report of increased VEGF protein expression by ~15% 2 h post resistance in lean individuals (Gavin et al, 2007). However, differences in analysis techniques could account for the variations, as Gavin and colleagues used ELISA for analysis which is thought to be more sensitive than Western Blot techniques (Kuijf et al,

2009). Further investigations are needed in order to develop an accurate time course of VEGF protein expression in the skeletal muscle following resistance exercise.

Increases in VEGF receptor mRNA have been observed at 4 h but not 2 h post resistance exercise in lean individuals (Gavin et al, 2007). Considering other factors involved in the present study, 3 h was chosen for the final biopsy with the hopes this would be sufficient time for changes in VEGF receptor mRNA to occur. The present study shows no increase in VEGF receptor mRNA in lean or obese individuals up to 3 h following resistance exercise. Similarly, exercise did not increase skeletal muscle VEGF receptor protein expression in lean or obese subjects. Gavin and colleagues also showed no changes in VEGF receptor protein expression 4 h post resistance exercise in lean individuals, as expected since protein changes occur after mRNA changes (Gavin et al, 2004). However, VEGF receptor protein expression increases have been observed 6 hours post aerobic exercise (Gustafsson et al, 2005). Since little is known about other factors involved in this study, a 6 h instead of 3 h post biopsy could have resulted in potentially missing important changes immediately following the acute resistance exercise bout. The last skeletal muscle biopsy was obtained 3 h post exercise in the present study. Therefore, 3 h may be insufficient to observe changes in VEGF receptor mRNA or protein expression following acute resistance exercise.

#### 4.3 Obesity and basal TSP-1 expression

TSP-1 is an angiostatic factor that is present in several tissues and is involved in cell adhesion, migration, and growth factor regulation (Lawler, 2002; Roberts, 1996; Sage & Bornstein, 1991). Comparisons of TSP-1 levels between lean and obese have

previously been investigated in the adipose tissue, as dynamic alterations in the extracellular matrix are required during the expansion of adipose tissue (Kong, Cavalera & Frangogiannis, 2014; Sun, Kusminski & Scherer, 2011). Vascularization and adipose tissue development are coupled from early development throughout adult life (Cranell, Hausman & Kral, 1997) Adipocytes play an important role in guiding and maintaining microvessel growth (Lau, 1990; Lau et al, 1990) and are a major site of TSP-1 secretion (Varma et al, 2008). Animal models demonstrate the lack of TSP-1 or its receptors as a protective mechanism against the development of obesity (Frazier et al, 2011; Kong et al, 2013; Maimaitiyiming et al, 2015). However, it has also been proposed that TSP-1 deficiency only improves glucose tolerance and increases insulin sensitivity with no effect on the development of obesity (Li et al, 2011). TSP-1 mRNA expression in the adipose tissue of human subjects suggests a positive correlation between TSP-1 expressed in the visceral adipose tissue and BMI (Matsuo et al, 2015; Varma et al, 2008). Evidence of increased TSP-1 expression in obese is also observed in other organs such as the heart (Gonzalez-Quesada et al, 2013) and blood vessels (Stenina et al, 2003).

To our knowledge, no previous studies have reported basal levels of TSP-1 expressed in the skeletal muscle of lean compared to obese individuals. Chronic delivery of TSP-1 to the skeletal muscle reduces capillarity, as shown in mouse models (Audet, Fulks, Stricker & Olfert, 2013). However, results of the present study show no differences in basal skeletal muscle TSP-1 mRNA despite significant differences in BMI. Therefore, another angiostatic factor may be responsible for the reduced capillary density present in the skeletal muscle of obese individuals.

#### 4.4 Obesity, exercise and TSP-1 expression

Investigations of TSP-1 mRNA expression following exercise have been reported with lean subjects. In Wistar rats, TSP-1 mRNA expression is increased 3.5 fold immediately following aerobic exercise, with further elevations 1 h post exercise before returning to baseline (Olfert, Breen, Gavin & Wagner, 2009). Three days of aerobic training attenuates this increase while the acute response is restored following 8 wks of training (Olfert, Breen, Gavin & Wagner, 2009). TSP-1 mRNA demonstrates a similar response to aerobic exercise in lean human subjects (Hoier et al, 2012). The current study is the first to investigate the TSP-1 mRNA and protein expression following acute resistance exercise in lean and obese subjects.

Aerobic exercise increases AMPK activity and TSP-1 expression at similar time points, within 2 h post exercise (Hoier et al, 2012; Olfert, Breen, Gavin & Wagner, 2009; Olfert et al, 2006; Olfert, Wagner, & Power, 2000; Sriwijitkamol et al, 2007; Stephens et al, 2002); however the present study shows no increase in AMPK phosphorylation or TSP-1 expression following resistance exercise. Though a negative finding, this is consistent with TSP-1 expression regulation by AMPK. Further investigation into skeletal muscle TSP-1 expression following exercise may be beneficial in order to investigate potential regulators of TSP-1 and establish an accurate time course of response.

#### 4.5 Obesity and basal p-AMPK and AMPK expression

The phosphorylation of AMPK regulates cellular energy usage by increasing the activity of energy conserving pathways while concurrently reducing the activity of

energy consuming pathways (Hawley et al, 1995). In obese Zucker rats, basal levels of skeletal muscle AMPK phosphorylation and the subsequent p-AMPK:AMPK ratio are significantly higher than their lean counterparts (Chen, Lee, Kuo & Ho, 2011). Other investigations, including the present study, have found no effect of obesity on basal AMPK activity (Lee-Young et al, 2011; Steinberg et al, 2004). Differences between human and animal models could in part account for the conflicting results. More investigation into the basal activity of AMPK between lean and obese, especially in human subjects, is needed.

#### 4.6 Obesity, exercise and p-AMPK and AMPK expression

AMPK is an important regulator of the body's response to stressors. One important stressor known to increase AMPK activity is exercise (Winder & Hardie, 1996). Mouse models have demonstrated that diet-induced obesity has a negative impact on AMPK activity in response to acute aerobic exercise (Lee-Young et al, 2011). Similar results have been shown in human studies involving moderate intensity aerobic exercise in obese subjects (Sriwijitkamol et al, 2007). In contrast, lean and obese Zucker rats show no difference in the activation of AMPK following 6 wks of aerobic exercise training (Chen, Lee, Kuo & Ho, 2011). However, obese Zucker rats have significantly lower AMPK phosphorylation following exercise training, while lean Zucker rats show no change (Chen, Lee, Kuo & Ho, 2011). This suggests aerobic exercise training could restore the AMPK response of obese subjects.

Investigations into AMPK activity in response to resistance exercise in lean individuals demonstrate conflicting results (Ahtiainen et al, 2015; Apro et al, 2015;



Dreyer et al, 2006; Silvennoinen et al, 2015). However, the phosphorylation of AMPK appears to be resistance exercise volume-dependent. Ahtiainen and colleagues observed an increased p-AMPK protein expression 30 min following 10 but not 5 sets of 10 RM using a leg press device (Ahtiainen et al, 2015). Previous reports suggest AMPK activity is increased within the first 30 min of exercise (Dreyer et al, 2006; Silvennoinen et al, 2015). The present study observed no changes in p-AMPK following 3 sets of knee extension. Therefore, the volume of the current exercise bout may have been insufficient to elicit a response since samples were obtained within the suggested time window.

#### 4.7 Limitations

As with all studies, limitations to the experimental design are present in the current report. Participants were both male and female and the menstrual cycle of female subjects were not controlled for. Hormonal differences could impact the skeletal muscle and therefore could result in increases variability within the results. Participants documented, but did not alter, their diet in this experiment. Subjects were asked to eat normally and record all food and beverage intake for three days preceding Day 2. Hours of sleep prior to Day 2 procedure were not controlled. At least two weeks separated Days 1 and 2; however the number of days between varied between subjects due to scheduling availability. Baseline biopsies were obtained from the non-working leg to avoid three biopsies on one single leg. Time of biopsies attempted to capture changes in muscle mRNA and proteins with limited information on their time courses following resistance exercise. Therefore, potential changes could have been missed. Due to limited information on the angiogenic response following resistance exercise, duration and intensity of the exercise bout may have been insufficient to produce changes.

#### 4.8 Future Directions

The increased VEGF expression reported in the current study further supports findings that resistance exercise training can elicit angiogenesis, though there is no difference between lean and obese individuals. However, the skeletal muscle of obese individuals is unable to maintain capillary density with muscle hypertrophy. It would be beneficial to study the long term angiogenic response to high volume resistance training in obese individuals, as there appears to be a disruption between the acute response and the development of capillaries. Other possible angiostatic factors that could be hindering capillary formation should be investigated.

## CHAPTER 5. CONCLUSION

In summary, the present study demonstrates that lean and obese individuals have a similar angiogenic response to acute resistance exercise. Future studies are needed in order to observe the effect of repeated bouts of resistance exercise on skeletal muscle angiogenesis between lean and obese individuals. Considerations into possible angiostatic factors other than TSP-1 contributing to reduced angiogenesis should be investigated.

## REFERENCES

## REFERENCES

- Abu-Farha M, Tiss A, Abubaker J, Khadir A, Al-Ghimlas F, Al-Khairi I, Baturcam E, Cherian P, Elkum N, Hammad M, John J, Kavalakatt S, Warsame S, Behbehani K, Dermime S, Dehbi M. Proteomics analysis of human obesity reveals the epigenetic factor HDAC4 as a potential target for obesity. *PLoS One*. 8(9):e75342, 2013.
- Adair TH, Montani JP. *Angiogenesis*. San Rafael (CA): Morgan & Claypool Life Sciences. Chapter 1, Overview of Angiogenesis, 2010.
- Ahtiainen JP, Walker S, Silvennoinen M, Kyrolainen H, Nindl BC, Hakkinen K, Nyman K, Selanne H, Hulmi JJ. Exercise type and volume alter signaling pathways regulating skeletal muscle glucose uptake and protein synthesis. *Eur J Appl Physiol*. 115:1835-1845, 2015.
- Amaral SL, Papanek PE, Greene AS. Angiotensin II and VEGF are involved in angiogenesis induced by short-term exercise training. *Am J Physiol Heart Circ Physiol*. 281:H1163-H1169, 2001.
- Andersen PO, Henriksson J. Capillary supply of the quadriceps femoris muscle of man: adaptive response to exercise. *J Physiol*. 270:677 -690, 1977.

Apro W, Moberg M, Hamilton DL, Ekblom B, van Hall G, Holmberg HC, Blomstrand E.

Resistance exercise-induced S6K1 kinase activity is not inhibited in human skeletal muscle despite prior activation of AMPK by high-intensity interval cycling. *Am J Physiol.* 308(6):E470-E481, 2015.

Audet GN, Fulks D, Stricker JC, Olfert IM. Chronic delivery of a thrombospondin-1 mimetic decreases skeletal muscle capillarity in mice. *PLoS One.* 8(2):e55953, 2013.

Bagavandoss P, Wilks JW. Specific inhibition of endothelial cell proliferation by thrombospondin. *Biochem Biophys Res Commun.* 170:867–872, 1990.

BenEzra D, Griffin BW, Maftzir G, Aharonov O. Thrombospondin and in vivo angiogenesis induced by basic fibroblast growth factor or lipopolysaccharide, *Invest Ophthalmol Vis Sci.* 34:3601-8, 1993.

Benoit H, Jordan M, Wagner H, Wagner PD. Effect of NO, vasodilator prostaglandins, and adenosine on skeletal muscle angiogenic growth factor gene expression. *J Appl Physiol.* 86:1512-1518, 1999.

Berggren JR, Boyle KE, Chapman WH, Houmard JA. Skeletal muscle lipid oxidation and obesity: influence of weight loss and exercise. *Am J Physiol Endocrinol Metab.* 294:E726-732, 2008.

Centers for Disease Control and Prevention. *National Diabetes Statistics Report: Estimates of Diabetes and Its Burden in the United States, 2014.* Atlanta: US Department of Health and Human Services, 2014.

Centers for Disease Control and Prevention. *Physical Activity and Health: The Benefits of Physical Activity.* Atlanta: US Department of Health and Human Service, 2015.

- Chen H, Herndon ME, Lawler J. The cell biology of thrombospondin-1. *Matrix Biology*. 19:597-614, 2000.
- Chen YC, Lee SD, Kuo CH, Ho LT. The effects of altitude training on the AMPK-related glucose transport pathway in the red skeletal muscle of both lean and obese Zucker rats. *High Alt Med Biol*. 12(4):371-378, 2011.
- Connolly DT, Heuvelman DM, Nelson R, Olander JV, Eppley BL, Delfino JJ, Siegel NR, Leimgruber RM, Feder J. Tumor vascular permeability factor stimulates EC growth and angiogenesis. *J Clin Invest*. 84:1470-1478, 1989.
- Costill D, Pearson D, Fink W. Impaired muscle glycogen storage after muscle biopsy. *J Appl Physiol*. 2245-2248, 1988.
- Crandall DL, Hausman GJ, Kral JG. A review of the microcirculation of adipose tissue: anatomic, metabolic, and angiogenic perspectives. *Microcirculation*. 4:211-232, 1997.
- Croley AN, Zwetsloot KA, Westerkamp LM, Ryan NA, AM, Hickner RC, Pofahl WE, Gavin TP. Lower capillarization, VEGF protein, and VEGF mRNA response to acute exercise in the vastus lateralis muscle of aged vs. young women. *J Appl Physiol*. 99:1872-1879, 2005.
- Dawson DW, Pearce SF, Zhong R, Silverstein RL, Frazier WA, Bouck NP. CD36 mediates the in vitro inhibitory effects of thrombospondin-1 on endothelial cells. *J Cell Biol*. 138(3):707-717, 1997.
- Degens H, Moore JA, Alway SE. Vascular endothelial growth factor, capillarization, and function of the rat plantaris muscle at the onset of hypertrophy. *Jpn J Physiol*. 53:181-191, 2003.



- Donnelly JE, Sharp T, Houmard J, Carlson MG, Hill JO, Whatley JE, Israel RG. Muscle hypertrophy with large-scale weight loss and resistance training. *Am J Clin Nutr.* 58(4):561-565, 1993.
- Doyle JL, Haas TL. The angiogenic response to skeletal muscle overload is not dependent on mast cell activation. *Microcirculation.* 17(7):548-556, 2010.
- Dreyer HC, Fuijita S, Cadenas JG, Chinkes DL, Volpi E, Rasmussen BB. Resistance exercise increases AMPK activity and reduces 4E-BP1 phosphorylation and protein synthesis in human skeletal muscle. *J Physiol.* 576(2):613-624, 2006.
- Dvorak HF, Orenstein NS, Carvalho AC, Churchill WH, Dvorak AM, Galli SJ, Feder J, Bitzer AM, Rypysc J, Giovenco P. Induction of a fibrin-gel investment: an early event in line 10 hepatocarcinoma growth mediated by tumor-secreted products. *J Immunol.* 122:166–174, 1979.
- Engerman RL, Pfaffenbach O, Davis MD. Cell turnover of capillaries. *Lab Invest.* 17:738-43, 1967
- Ferrara N, Davis-Smyth T. The biology of vascular endothelial growth factor. *Endocr Rev.* 18(1):4-25, 1997.
- Ferrara N, Henzel WJ. Pituitary follicular cells secrete a novel heparin-binding growth factor specific for vascular endothelial cells. *Biochem Biophys Res Commun.* 161:851-858, 1989.
- Ferrara N, Keyt B. Vascular endothelial growth factor: basic biology and clinical implications. *EXS.* 79:209-232, 1997.

- Frazier EF, Isenberg JS, Shiva S, Zhao L, Schlesinger P, Dimitry J, Abu-Asab MS, Tsokos M, Roberts DD, Frazier WA. Age-dependent regulation of skeletal muscle mitochondria by the thrombospondin-1 receptor CD47. *Matrix Biology*. 30(2):154-161, 2011.
- Fujii N, Hayashi T, Hirshman MF, Smith JT, Habinowski SA, Kaijser L, Mu J, Ljungqvist O, Birnbaum MJ, Witters LA, Thorell A, Goodyear LJ. Exercise induces isoform-specific increase in 5'AMP-activated protein kinase activity in human skeletal muscle. *Biochem Biophys Res Commun*. 273:1150-1155, 2000.
- Gavin TP. Basal and Exercise-Induced Regulation of Skeletal Muscle Capillarization. *Exercise and Sport Sciences Reviews*. 37:86-92, 2009.
- Gavin TP, Drew JL, Kubik CJ, Pofahl WE, Hickner RC. Acute resistance exercise increases skeletal muscle angiogenic growth factor expression. *Acta Physiol*. 191(2):139-146, 2007.
- Gavin TP, Robinson CB, Yeager RC, England JA, Nifong LW, Hickner RC. Angiogenic growth factor response to acute systemic exercise in human skeletal muscle. *J Appl Physiol*. 96:19-24, 2004.
- Gavin TP, Ruster RS, Carrithers JA, Zwetsloot KA, Kraus RM, Evans CA, Knapp DJ, Drew JL, McCartney JS, Garry JP, Hickner RC. No difference in the skeletal muscle angiogenic response to aerobic exercise training between young and aged men. *J Physiol*. 585(1):231-239, 2007.

- Gavin TP, Stallings HW, Zwetsloot KA, Westerkamp LM, Ryan NA, Moore RA, Pofahl WE, Hickner RC. Lower capillary density but no difference in VEGF expression in obese vs lean young skeletal muscle in humans. *J Appl Physiol.* 98(1):315-321, 2005.
- Geisler S, Brinkmann C, Schiffer T, Kreutz T, Bloch W, Brixius K. The influence of resistance training on patients with metabolic syndrome-significance of changes in muscle fiber size and muscle fiber distribution. *J Strength Cond Res.* 25(9):2598-2604, 2011.
- Gonzalez-Quesada C, Cavalera M, Biernacka A, Kong P, Lee DW, Saxena A, Frunza O, Dobaczewski M, Shinde AV, Frangogiannis NG. Thrombospondin-1 induction in the diabetic myocardium stabilizes the cardiac matrix while promoting vascular rarefaction through angiopoietin-2 upregulation. *Circ Res.* 113(12):1331-1344, 2013.
- Greenaway J, Lawler J, Moorehead R, Bornstein P, Lamarre J, Petrik J. Thrombospondin-1 inhibits VEGF levels in the ovary directly by binding and internalization via the low density lipoprotein receptor-related protein-1 (LRP-1). *J Cell Physiol.* 210:807-818, 2007.
- Gupta K, Gupta P, Wild R, Ramakrishnan S, Hebbel RP. Binding and displacement of vascular endothelial growth factor (VEGF) by thrombospondin: Effect on human microvascular endothelial cell proliferation and angiogenesis. *Angiogenesis.* 3:147-158, 1999.

- Gustafsson T, Puntchart A, Kaijser L, Jansson E, Sundberg CJ. Exercise-induced expression of angiogenesis-related transcription and growth factors in human skeletal muscle. *Am J Physiol.* 276:H679-H685, 1999.
- Gustafsson T, Ameln H, Fischer H, Sundberg CJ, Timmons JA, Jansson E. VEGF-A splice variants and related receptor expression in human skeletal muscle following submaximal exercise. *J Appl Physiol.* 98:2137-2146, 2005.
- Hardie DG. Minireview: the AMP-activated protein kinase cascade: the key sensor of cellular energy status. *Endocrinology.* 144:5179–5183, 2003.
- Hawley SA, Selbert MA, Goldstein EG, Edelman AM, Carling D, Hardie DG. 5'-AMP activates the AMP-activated protein kinase cascade, and Ca<sup>2+</sup>/calmodulin the calmodulin-dependent protein kinase I cascade, via three independent mechanisms. *J Biol Chem.* 270:27186-27191, 1995.
- Hellsten Y, Hoier B. Capillary growth in human skeletal muscle: physiological factors and the balance between pro-angiogenic and angiostatic factors. *Biochem Soc Trans.* 42:1616-1622, 2014.
- Hermansen L, Wachtlova M. Capillary density of skeletal muscle in well-trained and untrained men. *J Appl Physiol.* 30:860- 863, 1971.
- Hida K, Wada J, Zhang H, Hiragushi K, Tsuchiyama Y, Shikata K, Makino H: Identification of genes specifically expressed in the accumulated visceral adipose tissue of OLETF rats. *J Lipid Res.* 41:1615 –1622, 2000.
- Hoffner L, Nielsen JJ, Langberg H, and Hellsten Y. Exercise but not prostanoids enhance levels of vascular endothelial growth factor and other proliferative agents in human skeletal muscle interstitium. *J Physiol.* 550:217-225, 2003.

- Hoier B, Nordsborg N, Andersen S, Jensen L, Nybo L, Bangsbo J, Hellsten Y. Pro- and anti-angiogenic factors in human skeletal muscle in response to acute exercise and training. *J Physiol.* 590:595-606, 2011.
- Hoier B, Prats C, Qvortrup K, Pilegaard H, Bangsbo J, Hellsten Y. Subcellular localization and mechanism of secretion of vascular endothelial growth factor in human skeletal muscle. *FASEB J.* 27:3496-3504, 2012.
- Hoier B, Rufener N, Bojsen-Moller J, Bangsbo J, Hellsten Y. The effect of passive movement training on angiogenic factors and capillary growth in human skeletal muscle. *J Physiol.* 588:3833-3845, 2010.
- Holloszy JO, Coyle EF. Adaptations of skeletal muscle to endurance exercise and their metabolic consequences. *J Appl Physiol.* 56(4):831-838, 1984.
- Hong SB, Lee JJ, Kim SH, Suh YJ, Han JY, Kim YS, Nam M. The effects of adiponectin and inflammatory cytokines on diabetic vascular complications in obese and non-obese patients with type 2 diabetes mellitus. *Diabetes Res Clin Pract.* 111:58-65, 2015.
- Hudlicka O, Brown M, Egginton S. Angiogenesis in skeletal and cardiac muscle. *Physiol Rev.* 72:369-417, 1992.
- Huey KA, Smith SA, Sulaeman A, Breen EC. Skeletal myofiber VEGF is necessary for myogenic and contractile adaptations to functional overload of the plantaris in adult mice. *J Appl Physiol.* 120(2):188-195, 2016.
- Ikeda E, Achen MG, Breier G, Risau W. Hypoxia-induced transcriptional activation and increased mRNA stability of vascular endothelial growth factor in C6 glioma cells. *J Biol Chem.* 270:19761-19766, 1995.

- Inoue M, Jiang Y, Barnes II RH, Tokunaga M, Martinez-Santibanez G, Geletka L, Lumeng CN, Buchner DA, Chun TH. Thrombospondin 1 mediates high-fat diet-induced muscle fibrosis and insulin resistance in male mice. *Endocrinology*. 154(12):4548-4559, 2013.
- Iruela-Arispe ML, Bornstein P, Sage EH. Thrombospondin exerts an antiangiogenic effect on cord formation by endothelial cells in vitro. *Proc Natl Acad Sci USA*. 88:5026-5030, 1991.
- Iruela-Arispe ML, Porter P, Bornstein P, Sage EH. Thrombospondin-1, an inhibitor of angiogenesis is regulated by progesterone in the human endometrium. *J Clin Invest*. 97:403-412, 1995.
- Isenberg JS, Ridnour LA, Dimitry J, Frazier WA, Wink DA, Roberts DD. CD47 is necessary for inhibition of nitric oxide-stimulated vascular cell responses by thrombospondin-1. *J Biol Chem*. 281(36):26069-26080, 2006.
- Kalckar HM. The role of myokinase in transphosphorylations- The enzymatic action of myokinase on adenine nucleotides. *J Biol Chem*. 148:127-137, 1943.
- Katta A, Kakarla SK, Manne NDPK, Wu M, Kundla S, Kolli MB, Nalabotu SK, Blough ER. Diminished muscle growth in the obese Zucker rat following overload is associated with hyperphosphorylation of AMPK and dsRNA-dependent protein kinase. *J Appl Physiol*. 113(3):377-384, 1985.
- Keyt B, Nguyen H, Berleau L, Duarte C, Park J, Chen H, Ferrara N. Identification of VEGF determinants for binding Flt-1 and KDR receptors, generation of receptor-selective VEGF variants by site-directed mutagenesis. *J Biol Chem*. 271:5638-5646, 1996.

- Koch S, Tugues S, Li X, Gualandi L, Claesson-Welsh L. Signal transduction by vascular endothelial growth factor receptors. *Biochem.* 437(2):169-183, 2011.
- Kong P, Gonzalez-Quesada C, Li N, Cavalera M, Lee DW, Frangogiannis NG. Thrombospondin-1 regulates adiposity and metabolic dysfunction in diet-induced obesity enhancing adipose inflammation and stimulating adipocyte proliferation. *Am J Physiol.* 305(3):E439-E450, 2013.
- Kong P, Cavalera M, Frangogiannis NG. The role of thrombospondin (TSP)-1 in obesity and diabetes. *Adipocyte.* 3(1):81-84, 2014.
- Kraus RM, Stallings HW III, Yeager RC, Gavin TP. Circulating plasma VEGF response to exercise in sedentary and endurance-trained men. *J Appl Physiol.* 96:1445-1450, 2004.
- Kudo N, Barr AJ, Barr RL, Desai S, Lopaschuk GD. High rates of fatty acid oxidation during reperfusion of ischemic hearts are associated with a decrease in malonyl-CoA levels due to an increase in 5'-AMP-activated protein kinase inhibition of acetyl-CoA.. *J Biol Chem.* 270(29):17513-20, 1995.
- Kuijf ML, Eurelings M, Tio-Gillen AP, Van Doorn PA, Van Den Berg LH, Hooijkaas H, Stork J, Notermans NC, Jacobs BC. Detection of anti-MAG antibodies in polyneuropathy associated with IgM monoclonal gammopathy. *Neurology.* 73(9):688-695, 2009.
- Kuroki M, Voest EE, Amano S, Beerepoot LV, Takashima S, Tolentino M, Kim RY, Rohan RM, Colby KA, Yeo KT, Adamis AP. Reactive oxygen intermediates increase vascular endothelial growth factor expression in vitro and in vivo. *J Clin Invest.* 98:1667–1675, 1996.

- Lash, JM, Sherman WM, Hamlin RL. Capillary basement membrane thickness and capillary density in sedentary and trained obese Zucker rats. *Diabetes*. 38(7):854, 1989.
- Lau DC. Nature and nurture in adipocyte development and growth. *Int J Obes*. 14(3):153-157, 1990.
- Lau DC, Shillabeer G, Wong KL, Tough SC, Russell JC. Influence of paracrine factors on preadipocyte replication and differentiation. *Int J Obes*. 14(2):193-201, 1990.
- Lawler J. The functions of thrombospondin-1 and -2. *Curr Opin Cell Biol*. 12:634-640, 2000.
- Lawler J. The structural and functional properties of thrombospondin. *Blood*. 67:1197-1209, 1986.
- Lawler J. Thrombospondin-1 as an endogenous inhibitor of angiogenesis and tumor growth. *Cell Mol Med*. 6(1):1-12, 2002.
- Lawler JW, Slayter HS, Coligan JE. Isolation and characterization of a high molecular weight glycoprotein from human blood platelets. *J Biol Chem*. 253:8609-8616, 1978.
- Lawler PR, Lawler J. Molecular basis for the regulation of angiogenesis by thrombospondin-1 and -2. *Cold Spring Harb Perspect Med*. 2(5):a006627, 2012.
- Lee JM, Okumura MJ, Davis MM, Herman WH, Gurney JG. Prevalence and determinants of insulin resistance among US adolescents. *Diabetes Care*. 29(11):2427-2432, 2006.



- Lee-Young RS, Ayala JE, Fueger PT, Mayes WH, Kang L, Wasserman DH. Obesity impairs skeletal muscle AMPK signaling during exercise: role of AMPK $\alpha$ 2 in the regulation of exercise capacity *in vivo*. *Intl J Ob*. 35:982-989, 2011.
- Li Y, Tong X, Rumala C, Clemons K, Wang S. Thrombospondin1 deficiency reduces obesity-associated inflammation and improves insulin sensitivity in a diet-induced obese mouse model. *PLoS One*. 6(10):e26656, 2011.
- Lillioja S, Young AA, Culter CL, Ivy JL, Abbott WGH, Zawadzki JK, Yki-Jarvinen H, Christin L, Secomb TW, Bogardus C. Skeletal muscle capillary density and fiber type are possible determinants of in vivo insulin resistance in man. *J Clin Invest*. 80:415-424, 1987.
- Maimaitiyiming H, Norman H, Zhou Q, Wang S. CD47 deficiency protects mice from diet-induced obesity and improves whole body glucose tolerance and insulin sensitivity. *Sci Rep*. 5:8846, 2015.
- Maisonpierre PC, Suri C, Jones PF, et al. Angiopoietin-2, a natural antagonist for Tie2 that disrupts in vivo angiogenesis. *Science*. 277:55-61, 1997.
- Malek MH, Olfert IM. Global deletion of thrombospondin-1 increases cardiac and skeletal muscle capillarity and exercise capacity. *Exp Physiol*. 94:749-760, 2009.
- Marsin AS, Bertrand L, Rider MH, Deprez J, Beauloye C, Vincent MF, Van den Berghe G, Carling D, Hue L. Phosphorylation and activation of heart PFK-2 by AMPK has a role in the stimulation of glycolysis during ischemia. *Curr Biol*. 10:1247-1255, 2000.

- Matsuo Y, Tanaka M, Yamakage H, Sasaki Y, Muranaka K, Haha H, Ikai I, Shimatsu A, Inoue M, Chun TH, Satoh-Asahara N. Thrombospondin 1 as a novel biological marker of obesity and metabolic syndrome. *Metabolism*. 64(11):1490-1499, 2015.
- Matthews D, Hosker J, Rudenski A, Naylor B, Treacher D, Turner R. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*. 28(7):412-419, 1985.
- McCall GE, Byrnes WC, Dickinson A, Pattany PM, and Fleck SJ. Muscle fiber hypertrophy, hyperplasia, and capillary density in college men after resistance training. *J Appl Physiol*. 81:2004-2012, 1996.
- McKeown NM, Meigs JB, Liu S, Saltzman E, Wilson PWF, Jacques PF. Carbohydrate nutrition, insulin resistance, and the prevalence of the metabolic syndrome in the Framingham offspring cohort. *Diabetes Care*. 27(2):538-546, 2004.
- Millauer, B., S. Witzigmann-Voos, H. Schnurch, R. Martinez, N. P. H. Moiler, W. Risau, and A. Ullrich. High affinity VEGF binding and developmental expression suggest Flk-1 as a major regulator of vasculogenesis and angiogenesis. *Cell*. 72:835-846, 1993.
- Muniyappa R, Iantorno M, Quon MJ. An integrated view of insulin resistance and endothelial dysfunction. *Endocrinol Metab Clin North Am*. 37(3):685, 2008.
- Nicosia RF, Tuszynski GP. Matrix-bound thrombospondin promotes angiogenesis in vitro, *J. Cell Biol*. 124:183-93, 1994.
- Olfert IM, Breen EC, Gavin TP, Wagner PD. Temporal thrombospondin-1 mRNA response in skeletal muscle exposed to acute and chronic exercise. *Growth Factors*. 24(4):253-259, 2006.

Olfert IM, Howlett RA, Tang K, Dalton ND, Gu Y, Peterson KL, Wagner PD, Breen EC.

Muscle-specific VEGF deficiency greatly reduces exercise endurance in mice. *J Physiol.* 587:1755-1767, 2009.

Olfert IM, Wagner P, Power G. Chronic hypoxia, exercise training and skeletal muscle capillarity: angiogenic regulation and morphological consequences. *ProQuest Dissertations Publishing*, 2000.

Ouchi N, Shibata R, Walsh K. AMP-activated protein kinase signaling stimulates VEGF expression and angiogenesis in skeletal muscle. *Circ Res.* 96:838-846, 2005.

Pages G, Berra E, Milanini J, Levy AP, Pouyssegur J. Stress-activated protein kinases (JNK and p38/HOG) are essential for vascular endothelial growth factor mRNA stability. *J Biol Chem.* 275:26484–26491, 2000.

Parvaresh K, Huber A, Brochin R, Bacon P, McCall G, Huey K, Hyatt JP. Acute VEGF expression during hypertrophy is muscle phenotype-specific and localizes as a striated pattern within fibres. *Exp Physiol.* 95:1098-1106, 2010.

Paturi S, Gutta AK, Kakarla SK, Katta A, Arnold EC, Wu M, Rice KM, Blough ER. Impaired overload-induced hypertrophy in obese Zucker rat slow-twitch skeletal muscle. *J Appl Physiol.* 108(1):7-13, 2010.

Pollock ML, Vincent KR. Resistance training for health. *The President's Council on Physical Fitness and Sports Research Digest.* 2(8), 1996.

Primo L, Ferrandi C, Roca C, Marchio S, di Blasio L, Alessio M, Bussolino F.

Identification of CD36 molecular features required for its in vitro angiostatic activity. *FASEB J.* 19:1713-1715, 2005.

- Rasmussen BB, Winder WW. Effect of exercise intensity on skeletal muscle malonyl-CoA and acetyl-CoA carboxylase. *J Appl Physiol.* 83:1104–1109, 1997.
- Rattigan S, Bussey CT, Ross RM, Richards SM. Obesity, insulin resistance, and capillary recruitment. *Microcirculation.* 14:299-309, 2007.
- Roberts DD. Regulation of tumor growth and metastasis by thrombospondin-1. *FASEB J.* 10(10):1183-1191, 1996.
- Ryan NA, Zwetsloot KA, Westerkamp LM, Hickner RC, Pofahl WE, Gavin TP. Lower skeletal muscle capillarization and VEGF expression in aged vs. young men. *J Appl Physiol.* 100:178-185, 2006.
- Sage EH, Bornstein P. Extracellular proteins that modulate cell-matrix interactions. *J Biol Chem.* 266:14831-14834, 1991.
- Schalkwijk CG, Stehouwer CD. Vascular complications in diabetes mellitus: the role of endothelial dysfunction. *Clin Sci.* 109:143-159, 2005.
- Schantz P. Capillary supply in hypertrophied human skeletal muscle. *Acta Physiol Scand.* 114:635-637, 1982.
- Shono N, Mizuno M, Nishida H, Higaki Y, Urata H, Tanaka H, Quistorff B, Saltin B, Shindo M, Nishizumi M. Decreased skeletal muscle capillary density is related to higher serum levels of low-density lipoprotein cholesterol and apolipoprotein B in men. *Metabolism.* 48(10):1267-1271, 1999.
- Shweiki D, Itin A, Soffer D, Keshet E. Vascular endothelial growth factor induced by hypoxia may mediate hypoxia-initiated angiogenesis. *Nature.* 359:843-845, 1992.

- Silvennoinen M, Ahtiainen JP, Hulmi JJ, Pekkala S, Taipale RS, Nindl BC, Laine T, Hakkinen K, Selanne H, Kyrolainen H, Kainulainen H. PGC-1 isoforms and their target genes are expressed differently in human skeletal muscle following resistance and endurance exercise. *Physiol Rep*. 3:e12563, 2015.
- Sriwijitkamol A, Coletta DK, Wajcberg E, Balbontin GB, Reyna SM, Barrientes J, Eagan PA, Jenkinson CP, Cersosimo E, DeFronzo RA, Sakamoto K, Musi N. Effect of acute exercise on AMPK signaling in skeletal muscle of subjects with type 2 diabetes: a time-course and dose-response study. *Diabetes*. 56(3):836-848, 2007.
- Steinberg GR, Smith AC, Van Denderen BJ, Chen Z, Murthy S, Campbell DJ, Heigenhauser GJ, Dyck DJ, Kemp BE. AMP-activated protein kinase is not down-regulated in human skeletal muscle of obese females. *J Clin Endocrinol Metab*. 89(9):4575-4580, 2004.
- Stenina OI, Krukovets I, Wang K, Shou Z, Forudi F, Penn MS, Topol EJ, Plow EF. Increased expression of thrombospondin-1 in vessel wall of diabetic Zucker rat. *Circulation*. 107:3209-3215, 2003.
- Stephens TJ, Chen ZP, Canny BJ, Michell BJ, Kemp BE, McConnell GK. Progressive increase in human skeletal muscle AMPK  $\alpha$ 2 activity and ACC phosphorylation during exercise. *Am J Physiol*. 282:E688-E694, 2002.
- Sun K, Kusminski CM, Scherer PE. Adipose tissue remodeling and obesity. *J Clin Invest*. 121:2094-2101, 2011.
- Takahashi A, Kureishi Y, Yang J, Luo Z, Guo K, Mukhopadhyay D, Ivashchenko Y, Branellec D, Walsh K. Myogenic Akt signaling regulates blood vessel recruitment during myofiber growth. *Mol Cell Biol*. 22:4803-4814, 2002.

- Tang K, Breen EC, Gerber HP, Ferrara NM, Wagner PD. Capillary regression in vascular endothelial growth factor-deficient skeletal muscle. *Physiol Genomics*. 18:63-69, 2004.
- Tesch PA, Thorsson A, Kaiser P. Muscle capillary supply and fiber type characteristics in weight and power lifters. *J Appl Physiol*. 56:35- 38, 1984.
- Torgan CE, Brozinick JT, Kastello GM, Ivy JL. Muscle morphological and biochemical adaptations to training in obese Zucker rats. *J Appl Physiol*. 67(5):1807-1813, 1989.
- Trenerry MK, Carey KA, Ward AC, Cameron-Smith D. STAT3 signaling is activated in human skeletal muscle following acute resistance exercise. *J Appl Physiol*. 102(4):1483-1489, 2007.
- Varma V, Yao-Borengasser A, Bodles AM, Rasouli N, Phanavanh B, Nolen GT, Kern EM, Nagarajan R, Spencer HJ III, Lee MJ, Fried SK, McGehee RE Jr, Peterson CA, Kern PA. Thrombospondin-1 is an adipokine associated with obesity, adipose inflammation, and insulin resistance. *Diabetes*. 57(2):432-439, 2008.
- Voros G, Maquoi E, Demeulemeester D, Clerx N, Collen D, Lijnen HR: Modulation of angiogenesis during adipose tissue development in murine models of obesity. *Endocrinology*. 146:4545-4554, 2005.
- Wagner PD, Olfert IM, Tang K, Breen EC. Muscle-targeted deletion of VEGF and exercise capacity in mice. *Respir Physiol Neurobiol*. 151:159-166, 2006.
- Waltenberger J, Claesson-Welsh L, Siegbahn A, Shibuya M, Heldin CH. Different signal transduction properties of KDR and Flt1, two receptors for vascular endothelial growth factor. *J Biol Chem*. 269:26988-26995, 1994.

- Walton RG, Finlin BS, Mula J, Long DE, Zhu B, Fry CS, Westgate PM, Lee JD, Bennett T, Kern PA, Peterson CA. Insulin-resistant subjects have normal angiogenic response to aerobic exercise training in skeletal muscle, but not in adipose tissue. *Physiol Rep.* 3(6):e12415, 2015.
- Wang D, Huang HJ, Kazlauskas A, Cavenee WK. Induction of vascular endothelial growth factor expression in endothelial cells by platelet-derived growth factor through the activation of phosphatidylinositol 3-kinase. *Cancer Res.* 59:1464-1472, 1999.
- Winder WW, Hardie DG. Inactivation of acetyl-CoA carboxylase and activation of AMP-activated protein kinase in muscle during exercise. *Am J Physiol.* 270:E299-E304, 1996.
- Winzen R, Kracht M, Ritter B, Wilhelm A, Chen CY, Shyu AB, Muller M, Gaestel M, Resch K, Holtmann H. The p38 MAP kinase pathway signals for cytokine-induced mRNA stabilization via MAP kinase-activated protein kinase 2 and an AU-rich region-targeted mechanism. *EMBO J.* 18:4969-4980, 1999.
- Wojtaszewski JF, Nielsen P, Hansen BF, Richter EA, Kiens B. Isoform-specific and exercise intensity-dependent activation of 5'-AMP-activated protein kinase in human skeletal muscle. *J Physiol.* 528(1):221-226, 2000.
- Yamaguchi TP, Dumont DJ, Conlon RA, Breitman ML, Rossant J. Flk-1, an Flt-related receptor tyrosine kinase, is an early marker for endothelial cell precursors. *Development.* 118:489-498, 1993.

- Yamashita H, Sato N, Yamamoto M, Gasa S, Izawa T, Komabayashi T, Ishikawa M, Sata Y, Ohno H. Effect of endurance training on angiogenic activity in skeletal muscles. *Eur J Physiol.* 422:332-338, 1993.
- Zhu W, Smart EJ. Myristic acid stimulates endothelial nitric-oxide synthase in a CD36- and AMP kinase-dependent manner. *J Biol Chem.* 280:29543-29550, 2005.
- Ziche M, Morbidelli L, Masini E, Amerini S, Granger HJ, Maggi CA, Geppetti P, Ledda F. Nitric oxide mediates angiogenesis in vivo and endothelial cell growth and migration in vitro promoted by substance P. *J Clin Invest.* 94:2036-2044, 1994.
- Zwetsloot KA, Westerkamp LM, Holmes BF, Gavin TP. AMPK regulates basal skeletal muscle capillarization and VEGF expression, but is not necessary for the angiogenic response to exercise. *J Physiol.* 586(24):6021-6035, 2008.